Int. Agrophys., 2025, 39, 399-411 doi: 10.31545/intagr/204851

Morphophysiological adjustments and antioxidant defense in main tree species of southern China to simulated acid deposition**

Matoor M. Gilani^{1,2}, Ali Noman³, Taimoor H. Farooq⁴, Ghulam Yasin², Muhammad H.U. Rashid⁵, Muhammad Zubair², Muhammad W.K. Tarin⁶, Fahad M. Alzuaibr⁷, Bo Liu⁸, Xiangqing Ma¹*

¹Forestry College, Fujian Agriculture and Forestry University Fuzhou, 350002 Fujian Province, PR China ²Department of Forestry and Range Management, Bahauddin Zakariya University, Multan, 66000, Pakistan ³Department of Botany, Government College University Faisalabad, 38040, Pakistan

⁴Bangor College China, a joint unit of Bangor University and Central South University of Forestry and Technology Changsha 410004, China

⁵Department of Forestry and Range Management, Bahauddin Zakariya University, 38000, Pakistan

⁶Department of Forestry and Range Management, Agriculture University Dera Ismail Khan, 29111, Pakistan

⁷Department of Biology, Faculty of Science, University of Tabuk, Tabuk, 47713, Saudi Arabia

8 School of Life Sciences, Qufu Normal University, Qufu 273165, Shandong Province, China

Received February 26, 2025; accepted May 9, 2025

Abstract. Acid deposition has become a significant issue due to increased pollution and forest degradation. To examine the response of acid deposition in Fuzhou, a city in Fujian Province, we investigated the physiology, morphology, osmotic adaptation, photosynthetic characteristics, and antioxidant defense of Pinus massoniana, Cunninghamia lanceolata, Liquidambar formosana, Schima superba, and Phoebe zhennan. Seedlings were subjected to pH levels of 2.5, 3.5, 4.5, and 5.5 in a greenhouse experiment. Plant height, diameter, and biomass increased in the control group (6.5); however, pH 4.5 and 5.5 resulted in tall and thin seedlings. Height and diameter decreased under increased acid deposition specifically at pH 2.5 and 3.5. Biochemical parameters and chlorophyll contents were reduced under low pH levels. The activities of catalase, peroxidase, superoxide dismutase, malonaldehyde, and proline increased initially under low pH, but decreased later due to high acidity, whereas soluble sugar and protein contents were inhibited under higher acidity levels (pH 2.5 and 3.5). The findings showed that the seedlings responded differently but more sensitively to acid deposition below pH 4.5. The exposure to less acidic deposition had a slight effect on the development of these tree seedlings. This trial aids in detecting bioindicators and resilient species to acid deposition existing in forests and urban areas.

K e y w o r d s: acid rain, photosynthesis, antioxidant enzyme, gas exchange, osmotic adjustment

1. INTRODUCTION

Acid deposition is recognized as one of the most serious environmental concerns (Gilani et al., 2021). It causes significant damage to forest ecosystems in Scandinavian countries, the United States, Canada, and Europe (Grennfelt et al., 2020). To date Europe, northeast America and China have emerged as the three leading regions in the world impacted by acid deposition (Liu et al., 2022). Reports indicate that the losses caused by acid rain are alarming in several provinces of southern China (Chen et al., 2010). For instance, Sichuan province in southern China is being badly stricken by acid deposition (Zhang et al., 2021b). The pH of rainfall in southern China ranks among the lowest globally, placing the region third in terms of acid rain severity (Ren et al., 2024). Coastal cities are a major source of acid rain due to air pollution caused by industries, vehicles, and construction activities (Meng et al., 2019). This issue is predominantly caused by the use of coal for energy production culminating at addition of sulfate (SO_4^{2-}) (Cape, 1993). Fuzhou, the capital of Fujian Province, is substantially affected by acid deposition, with higher severity observed in spring and winter compared

^{© 2025} Institute of Agrophysics, Polish Academy of Sciences



^{*}Corresponding author e-mail: lxymxq@126.com

^{**}This work was supported by Cross-Strait Collaborative Innovation Center of Soil and Water Conservation (K80ND8003) (2015-2027).

to autumn and summer, particularly in the southeastern, southwestern, western, and northwestern areas (Zheng *et al.*, 2014). The acid deposition affected area has been reduced in recent years; however, this issue continues to restrict social development and economic growth of China (Ren *et al.*, 2024). Despite the growing concern about acid deposition, research on its ecological consequences has lagged in China (Chang *et al.*, 2022).

Environmental and ecosystem-level problems caused by both biotic and abiotic factors can disrupt plants in multiple ways (Yasin *et al.*, 2023). Plants detect and respond to these changes by using defense mechanisms for their survival (Hasan *et al.*, 2022). For example, tree leaching can alter nutrient cycling processes in forest ecosystems, resulting in physiological stress (Farooq *et al.*, 2020). Similarly, acid deposition exacerbates the loss of foliage nutrients, thereby reducing tree biomass as well as vertical and radial growth (Fan and Wang, 2000). As a result, leaves often become desiccated and die, leading to a decline in both aboveground and belowground biomass (Huang *et al.*, 2012). Studies have shown that acid deposition affects the process of photosynthesis (Dong *et al.*, 2017).

This photosynthesis is eventually impacted by acid deposition due to destruction of chloroplast structure (Sun et al., 2012) and a decrease in the chlorophyll content (Du et al., 2017). Additionally, acid deposition can damage the leaf cuticle *e.g.* through the removal of wax layers, thereby exacerbating plant injury and disrupting metabolic processes (Diatta et al., 2021). This also enhances reactive oxygen species (ROS) and increases the malondialdehyde (MDA) content (Chen et al., 2006). The activity of antioxidant enzymes, such as superoxide dismutase, peroxidase, and catalase, initially rises in response to oxidative stress but may eventually decline under higher acidity, weakening the plant's ability to resist ROS (Rahman et al., 2021). Nevertheless, the enzyme activity lowers under increased acidity, decreasing plant resistance to stress (Ma et al., 2020). Previous findings have shown that acid deposition significantly impacted the physiology and antioxidant defense mechanisms of various tree seedlings, including L. formosana (Feng et al., 2002) and P. massoniana (Liu et al., 2018). Exposure to acid deposition particularly at pH 3.5 and below can result in noticeable phenotypic differences in the leaves of both broadleaved and coniferous tree species due to physiological damage, ultimately affecting the overall health of forest ecosystems (Chen et al., 2014). Similarly, another study revealed that the conifers forests are more vulnerable to acid deposition than the evergreen broadleaf forests (Hu et al., 2014). Acid deposition disrupts the enzymatic and non-enzymatic defense within plant cells, intensifying oxidative stress (Debnath et al., 2021). This oxidative stress can cause detrimental effects, such as reduced biomass and chlorophyll content as well as increased leaf dryness and mortality (Wei et al., 2017). Another study highlighted that populations of certain forest species have already declined due to acid deposition (Li *et al.*, 2013). The levels of soluble proteins, sugars, and proline are important indicators of stress resistance under environmental hazards and are directly influenced by acid deposition (Hua *et al.*, 2022). Proline accumulation is a key stress indicator in response to pathogens, heavy metals (Yasin *et al.*, 2021), air pollution, nutrient deficiency (Noman *et al.*, 2018), and increased sugar levels (Naheed *et al.*, 2021). This study aimed to investigate the responses of trees under elevated acid deposition. A previous study showed that both conifers and broadleaved species showed structural damage after exposure to acid rain (Cape, 1993).

Chinese fir (Cunninghamia lanceolata) is a native species that accounts for approximately 25% of plantations in southern China (Farooq et al., 2019). Previous findings revealed that this conifer species was damaged by acid rain in seven provinces of southern China (Blanco et al., 2012). Masson pine (Pinus massoniana) is another indigenous species in southern China which is characterized by its durability and fast growth and is a pioneer species for reforestation and afforestation (Zhou et al., 2020). Reports have indicated that *P. massoniana* is sensitive to acid deposition, with yield reductions of up to 43% when exposed to rain with a pH below 4.0 (Wu Gang, 1998; Farooq et al., 2021). Phoebe zhennan is a valuable species in China, prized for its highly durable wood, commonly known as "golden thread nanmu"(Zhu et al., 2022). As a result, the plantation area for this species has expanded rapidly. However, poor management practices have limited its production potential (Cheng et al., 2023). Liquidambar formosana is one of the ideal species for tree adjustment in plantations, and it is frequently chosen as an urban tree species (Wu et al., 2022). Several counties in Fujian Province have selected this species for experimental trials to enhance survival and germination rates (Gilani et al., 2023). S. superba is an evergreen broadleaf species found in subtropical forests in China (Chen et al., 2021). Earlier studies reported that its population size has declined throughout the last decade (Chen et al., 2013). Still, it is not evident whether its growth is influenced by acid deposition or by leaf injury (Yao et al., 2016). It was found that the population of S. superba and particularly L. formosana declined during the last few years (Liu et al., 2007). We hypothesized that acid rain can differentially affect photosynthetic attributes, lipid peroxidation, and antioxidant enzyme activity, thereby reducing biomass production in broadleaved species, pines, and conifers. To test this hypothesis, we conducted experiments with the following objectives: 1) to assess changes in gas exchange parameters under acid deposition; 2) to estimate the comparative activity of antioxidant enzymes and the extent of membrane damage due to acid rain; and 3) to record comparative biomass production in broadleaved species, pines, and conifers affected by acid rain. This trial aims to provide insight into the mechanisms by which acid deposition



Fig. 1. Research layout.

affects these species (Fig. 1). We anticipate that the findings will have practical implications for the protection and management of forests exposed to acid deposition.

2. MATERIALS AND METHODS

2.1. Experimental material

A one-year pot experiment was carried out in the green house of Fujian Agriculture and Forestry University, Fuzhou, Fujian. The area lies in a subtropical region with high temperatures in summer. It also encounters typhoons originating from Pacific storms. The average annual deposition is 1400-2000 mm. One-year-old seedlings of *P. massoniana, C. lanceolata, L. formosana, S. superba,* and *P. zhennan* were shifted into 125 pots and arranged in a completely randomized design (CRD) with five replications for each treatment and 25 plants per species. Pots of a uniform size of 20 cm from the top and 15.5 cm from the bottom with a height of 16.5 cm) were established in a greenhouse. The seedlings were grown in commercial soil containing organic matter.

2.2. Acid rain solutions

The acid solution was prepared with $1 \text{ N H}_2\text{SO}_4$ and 1 N HNO₃ in the ratio of 10 to 1 by chemical equivalents, diluted with distilled water to attain solutions with pH values of 2.5, 3.5, 4.5, and 5.5. 500 ml of the solution was given to each pot once a week. This was determined according to the

composition of rainfall by (Fan and Wang, 2000). Distilled water was used as the (CK) control group (pH 6.5). The seedlings were placed in a greenhouse which was covered with shade cloth to protect it from extreme temperature. The one-year trial began in August. The spray method was uniformly used throughout the experiment. The spray was applied at the top of the plants to distinguish the degree of stress due to different acidities. The sprayer contained acid resistant materials, the nozzle was made of stainless steel, and the water pipe was made of plastic.

2.3. Growth parameters

Seedling height (cm) was determined from the base to the top using a tape, and root collar diameter (mm) was determined using a vernier caliper (Mofunanya and Soonen, 2017). The seedlings were harvested and washed carefully for biomass measurement. Later, they were oven-dried at 65°C for 48 h. At the time of harvesting, the root/shoot ratio was measured by dividing the root with shoot of the plants.

2.4. Gas exchange measurements

Measurements were taken between 9:00-11:00 a.m. to ensure consistency. All measurements were recorded using a photosynthesis measurement system (LI-6400) at 1000 μ mol m⁻² s⁻¹ PPFD. Three similar leaves were

chosen from the top of each seedling. We obtained three measurements and used the average as the result. The measurement included the transpiration rate (T_r) , intercellular concentration (C_i) , photosynthetic rate (P_n) , and stomatal conductance (G_s) .

2.5. Determination of biochemical and photosynthetic pigments

The concentrations of photosynthetic pigments (chlorophyll *a*, chlorophyll *b*, and carotenoids) in leaf tissues were determined according to the methods described by (Lichtenthaler and Wellburn, 1983). Seedlings were either instantly used for testing or frozen in liquid nitrogen at -80°C. The pigment concentrations were determined using the following equations, based on the absorbance of the extract at 663, 645, and 470 nm, and expressed as mg g⁻¹:

$$Chlorophyll\ a\ content = \frac{(12.21D663 - 2.81D645)V}{1000W},\qquad(1)$$

$$Chlorophyll \ b \ content = \frac{(20.13D645 - 5.03D663)V}{1000W}, \qquad (2)$$

$$Carotenoids = \frac{(1000D470 - 3.27C_a - 104C_b)}{229} \frac{V}{1000W}, \qquad (3)$$

where: D is the absorbance of the supernatant at the specified wavelengths (nm), V is the extract volume, W is the sample weight, C_a represents chlorophyll a, C_b represents chlorophyll b.

2.6. Lipid peroxidation and enzymatic antioxidant measurement

The ELISA kit was used as stated by (Shanghai Enzymelinked Biotechnology Co. Ltd). SOD was observed using Nitro blue tetrazolium chloride, POD with guaiacol, and CAT with UV-Vis Spectroscopy (Ahmed *et al.*, 2013). The colorimetric method was employed to measure MDA contents (Hodges *et al.*, 1999).

2.7. Total soluble protein and sugars

Soluble protein was measured with an earlier technique proposed by Bradford (1976), while soluble sugar was determined according to the protocol used by Yemm and Willis (1954). Proline in leaves was identified at 520 nm (Bates *et al.*, 1973).

2.8. Statistical analysis

All data were analyzed using SPSS 20.0 and Statistics 8.1, while all figures were drawn by GraphPad Prism 8 Software. Two-way analysis was used and significance of differences between means was determined by applying Tukey HSD test at 5.

3. RESULTS

3.1. Growth parameters

The growth parameters, such as plant height and root collar diameter, decreased among all species and treatments. For example, the plant height in C. lanceolata decreased by 26.94, 21.26, 16.84, and 12.02% at pH 2.5, 3.5, 4.5, and 5.5. A more pronounced reduction in the plant height was observed in Schima superba, which showed decreases of 32.61, 27.63, 23.33, and 19.07% at the same pH levels, compared to the control group. Similarly, the root collar diameter decreased at all the pH levels, relative to the control. However, a greater reduction in the plant height was observed for S. superba at all the pH levels, which decreased by 32.61, 27.63, 23.33, and 19.07% at pH 2.5, 3.5, 4.5, and 5.5, respectively, compared to the control group. The root collar diameter decreased at all the pH levels, compared to the control; however, this reduction was higher at pH 2.5 in C. lanceolata, L. formosana, P. zhennan, and S. superba, compared to pH 3.5, pH 4.5, and pH 5.5, respectively (Table 1).

The leaf biomass in *C. lanceolata, P. massoniana, P. zhennan*, and *S. superba* decreased at pH 2.5 3.5, and 4.5; however, it increased in the studied species at pH 5.5.

Table 1. Effect of acid stress on plant height and root collar diameter of analyzed species (mean \pm SE)

| Tree parameter | Treatment | C. lanceolata | L. formosana | P. massoniana | P. zhennan | S. superba |
|------------------------------|-----------|---------------|--------------|---------------|--------------|--------------|
| Plant height (cm) | CK | 45.24±3.67a | 65.87±4.63a | 41.28±4.01a | 56.71±4.48a | 48.38±3.73a |
| | 2.5 | 33.05±1.69c | 43.55±2.21c | 23.48±1.62c | 34.18±1.89c | 32.60±2.17c |
| | 3.5 | 35.62±1.89bc | 45.33±3.30bc | 25.41±1.69c | 37.23±2.21c | 35.01±2.51bc |
| | 4.5 | 37.62±1.86b | 48.37±3.37b | 28.44±1.91bc | 40.92±2.44bc | 37.09±2.43b |
| | 5.5 | 39.80±2.31ab | 50.87±3.52b | 33.37±2.77bc | 44.74±2.69bc | 39.15±2.90ab |
| Root collar diameter (mm) | СК | 4.29±0.18a | 5.10±0.23a | 2.90±0.17a | 4.96±0.47a | 4.41±0.31a |
| | 2.5 | 3.56±0.06c | 4.24±0.06c | 2.48±0.04c | 3.74±0.26c | 3.17±0.17c |
| | 3.5 | 3.63±0.08c | 4.47±0.10bc | 2.44±0.08c | 4.34±0.30bc | 3.74±0.16bc |
| | 4.5 | 3.73±0.08bc | 4.81±0.22b | 2.75±0.10b | 4.44±0.30bc | 3.83±0.26bc |
| | 5.5 | 3.83±0.12b | 4.85±0.23ab | 2.52±0.11bc | 4.64±0.36a | 3.86±0.29b |

| Tree parameter | Treatment | C. lanceolata | L. formosana | P. massoniana | P. zhennan | S. superba |
|---------------------|-----------|------------------|------------------|---------------|-------------|------------|
| Leaf (g) | СК | 17.35±1.15a | 5.78±0.99a | 16.35±1.25a | 7.89±0.61b | 7.45±0.48a |
| | 2.5 | 9.12±0.35b | 2.61±0.28b | 9.32±0.39c | 6.01±0.06b | 4.03±0.48c |
| | 3.5 | 13.47±1.14b | 2.64±0.47b | 14.47±1.17b | 7.67±0.36b | 5.43±0.74b |
| | 4.5 | 10.59±0.67b | 3.01±0.29b | 11.59±0.69b | 5.59±0.71c | 5.11±0.29b |
| | 5.5 | 17.10±1.18a | 2.95±0.39b | 18.10±1.28a | 8.29±0.64a | 6.68±0.67b |
| | СК | 10.92±0.63a | 7.41±0.51a | 11.92±0.69a | 8.38±0.37a | 7.02±0.34a |
| | 2.5 | 6.00±0.43b | 5.71±1.06c | 6.10±0.49b | 5.89±0.61c | 5.08±0.53b |
| Stem (g) | 3.5 | 6.67±0.38b | 6.64±0.24b | 6.74±0.42b | 6.94±0.24b | 4.52±0.36c |
| | 4.5 | 7.71±0.71b | 6.69±0.28b | 7.73±0.76b | 7.96±0.71b | 5.47±0.22b |
| | 5.5 | 10.21±0.60a | 6.82±0.96b | 10.36±0.68a | 7.76±0.47b | 5.53±0.18b |
| | СК | 9.13±0.17a | 6.66±0.41a | 9.36±0.18a | 12.93±1.55a | 7.98±0.69a |
| | 2.5 | 6.35±0.70b | 3.46±0.08c | 6.39±0.71c | 8.69±0.61b | 4.78±0.25c |
| Root (g) | 3.5 | 7.13±0.60b | 5.95±0.32b | 7.16±0.61b | 9.31±0.59b | 5.53±0.43b |
| | 4.5 | 8.13±0.65b | 5.49±0.44b | 8.16±0.68b | 10.26±0.84b | 6.15±0.65a |
| | 5.5 | 7.98±0.89b | 6.62±1.40b | 7.87±0.85b | 12.57±0.81a | 6.07±0.52a |
| Root/Shoot Ratio | СК | 0.83±0.11b | 0.89±0.13ab | 0.78±0.09b | 1.54±0.23b | 1.13±0.25b |
| | 2.5 | 1.05±0.06a | 0.60±0.09c | 1.04±0.14a | 1.47±0.31b | 0.94±0.20c |
| | 3.5 | 1.06±0.09a | 0.89±0.06ab | 1.06±0.19a | 1.34±0.19c | 1.22±0.09a |
| | 4.5 | 1.05±0.23a | 0.82±0.12b | 1.14±0.08a | 1.28±0.11c | 1.12±0.13b |
| | 5.5 | $0.78{\pm}0.08b$ | $0.97{\pm}0.27a$ | 0.75±0.11b | 1.61±0.29a | 1.09±0.16b |

Table 2. Leaf, stem, root biomass and root/shoot ratio of analyzed species in response to acid stress (mean \pm SE)

In the case of L. formosana, the maximum leaf biomass (5.78 g) was computed in the control, and it decreased at all the pH levels with a minimum (2.61 g) at pH 2.5 (Table 2). The stem biomass decreased across all the species at all the pH levels, except at pH 5.5. The root biomass also declined in all the species under acid treatments, except at pH 4.5, where Cunninghamia lanceolata, Pinus massoniana, and Phoebe zhennan exhibited an increase, compared to the control group. The root/shoot ratio in C. lanceolata, L. formosana, and P. massoniana increased at pH 2.5, 3.5, and 4.5, except pH 5.5, where it decreased by 6.02% in C. lanceolata, 8.98% in L. formosana, and 3.84% in P. massoniana, respectively. The root/shoot ratio in P. zhennan decreased in all the treatments except pH 5.5, where it increased by 4.54%. Similarly, the root/shoot ratio in S. superba also showed a decreasing trend under all the pH treatments except pH 3.5, where it increased by 7.96%, compared to the control (Table 2).

3.2. Gas exchange measurements

The photosynthetic rate (P_n) decreased across all the treatments in all the studied species except *S. superba*. The P_n in *S. superba* increased by 9.82, 25.21, 26.41, and 45.4% at pH 2.5, 3.5, 4.5, and 5.5, respectively. Stomatal conductance (G_s) in *Cunninghamia lanceolata* increased under all the treatments except at pH 5.5, where it decreased by 16.03%, compared to the control. The G_s in *P. massonia*-

na, L. formosana, and P. zhennan was reduced at all the pH levels except pH 5.5, where it increased by 4.34% in P. massoniana, 10.97% in L. formosana, and 61.11% in *P. zhennan.* However, the G_s in *S. superba* showed an increasing trend at all the pH levels, compared to the control treatment. The intercellular CO_2 concentration (C_i) in C. lanceolata and P. massoniana increased across all pH levels. However, in L. formosana, Ci decreased by 15.37 and 41.1% at pH 2.5 and pH 3.5, respectively, while it increased by 1.81 and 36.23% at pH 4.5 and pH 5.5. Similarly, the C_i in S. superba increased by 17.95% at pH 3.5, compared with the control treatment. The transpiration rate (T_r) in C. lanceolata increased at all the pH levels except pH 5.5, where it decreased by 18.39%. The T_r in *P. massoniana*, L. formosana, and P. zhennan were lower at pH 2.5, 3.5, and 4.5; however, higher T_r was observed at pH 5.5. In the case of S. superba, a greater transpiration rate was observed at all the pH levels, compared with the control (Fig. 2).

3.3. Determination of biochemical and photosynthetic pigments

The chlorophyll contents in all the species decreased at all the pH levels (Table 3). The level of *chl a* in *C. lanceolata* decreased by 65.5, 58.2, 46.6, and 26.7% at pH 2.5, 3.5, 4.5, and 5.5. In the case of *S. superba*, it decreased by 64.25, 64.73, 71.49, and 68.11% at pH 2.5, 3.5, 4.5, and 5.5, respectively, compared to the control treatment. Similarly,



Fig. 2. a) Response of net photosynthetic rate (P_n) of photosynthetic photon flux (PPFD); b) stomatal conductance (G_s) to H_2O ; c) intercellular (C_i) CO₂ concentration; and d) transpiration rate (T_r) to treatment with simulated acid deposition.

| Tree parameter | Treatment | C. lanceolata | L. formosana | P. masoonian | P. zhennan | S. superba |
|--------------------------------------|-----------|--------------------|------------------|---------------------|-------------------|-------------------|
| Chl $a (mg g^{-1})$ | СК | 2.32±0.25a | 1.99±0.13a | 1.74±0.08a | 1.93±0.10a | 2.07±0.13a |
| | 2.5 | $0.80{\pm}0.07c$ | 0.83±0.05c | $0.43{\pm}0.04c$ | $0.97{\pm}0.03c$ | $0.74 \pm 0.06b$ |
| | 3.5 | $0.97{\pm}0.09c$ | 1.15±0.02b | 0.56±0.01b | $0.97{\pm}0.08c$ | 0.73±0.13b |
| | 4.5 | $1.24{\pm}0.04b$ | 1.52±0.03b | $0.71 {\pm} 0.05 b$ | $1.34{\pm}0.07b$ | 0.59±0.10b |
| | 5.5 | 1.70±0.04b | 1.76±0.05b | 0.99±0.11b | $1.21 \pm 0.07b$ | $0.66 \pm 0.09 b$ |
| | | | | | | |
| Chl $b (mg g^{-1})$ | CK | 1.69±0.04a | 1.62±0.12a | 1.27±0.05a | 1.23±0.12a | 1.55±0.11a |
| | 2.5 | $0.41 {\pm} 0.06b$ | 0.35±0.03b | 0.10±0.01c | $0.32{\pm}0.05b$ | $0.32 \pm 0.03 b$ |
| | 3.5 | $0.39{\pm}0.04b$ | 0.46 ± 0.06 | $0.26 \pm 0.008 b$ | $0.22 \pm 0.05c$ | $0.27 \pm 0.05 b$ |
| | 4.5 | $0.58{\pm}0.06b$ | 0.30±0.05c | $0.26 \pm 0.02b$ | 0.21±0.06c | 0.25±0.01c |
| | 5.5 | $0.48 \pm 0.06b$ | $0.42 \pm 0.06b$ | $0.24{\pm}0.02b$ | $0.48 \pm 0.07 b$ | $0.40 \pm 0.07 b$ |
| | | | | | | |
| Carotenoids (mg g ⁻¹) | CK | 0.63±0.15a | 0.89±0.12a | 0.81±0.03a | $0.62{\pm}0.09a$ | 0.94±0.05a |
| | 2.5 | $0.14{\pm}0.01c$ | $0.17 \pm 0.02c$ | 0.12±0.01c | 0.27 ± 0.02 c | $0.22 \pm 0.02b$ |
| | 3.5 | $0.21 \pm 0.04b$ | $0.25 \pm 0.02c$ | $0.13{\pm}0.004c$ | $0.30 \pm 0.04 b$ | $0.20 \pm 0.04 b$ |
| | 4.5 | $0.24{\pm}0.02b$ | 0.45±0.03b | $0.18 \pm 0.02b$ | 0.45±0.03b | 0.13±0.03c |
| | 5.5 | 0.44±0.03a | 0.50±0.04b | 0.28±0.05b | 0.28±0.03c | 0.56±0.04b |

Table 3. Chlorophyll *a*, chlorophyll *b*, and carotenoid content in response to acid stress (mean \pm SE)

NA

the *chl b* content in all the studied species decreased at all the pH levels. Likewise, the carotenoid contents under acid deposition declined with the increase in acidity. The carotenoid contents decreased by 77.8, 66.7, 61.9, and 30.2% in *C. lanceolata* and by 76.59, 78.72, 86.17, and 40.42% in *S. superba* at pH 2.5, 3.5, 4.5, and 5.5, respectively, as presented in Table 3.

The influence of acid deposition on enzymatic antioxidants in all the studied species is presented in Fig. 3. The findings revealed that all the enzymatic activities in the seedlings of studied species decreased with the increase in acidity levels. The CAT activity in C. lanceolata and L. formosana increased at all the pH levels. In the case of P. zhennan, the CAT activity decreased across all the pH levels except pH 2.5 where it increased by 19.08%, compared to the control. However, CAT in S. superba increased by 11.97, 19.74, 31.6, and 28.03% at pH 2.5, 3.5, 4.5, and 5.5, respectively. Similar to CAT, the SOD activity also showed a decreasing trend with the increase in acidity among all the species. The lowest SOD values were computed in the control treatment, whereas the highest SOD activity was observed at pH 5.5. The POD value increased by 0.43 and 1.63% at pH 2.5 and 3.5 in P. massoniana but decreased by 50.53 and 67.85% at pH 4.5 and pH 5.5, respectively.

Similarly, POD increased in all the other species at all the pH levels, compared to the control treatment, showing maximum POD values at pH 5.5 and minimum levels in the control group (Fig. 3). The MDA contents showed no significant variation in all the species under all the treatments. The MDA activity in *C. lanceolata, P. massoniana, L. formosana, P. zhennan*, and *S. superba* decreased at all the pH levels except pH 2.5, where it increased by 13.86, 13.26, 6.21, 4.24, and 8.3%, respectively.

In *C. lanceolata*, the soluble sugar content decreased by 44.25, 47.16 31.21, and 44.76% at pH 2.5, 3.5, 4.5, and 5.5, respectively, while the soluble sugar content in *S. superba* decreased at all the pH levels except pH 4.5, where it increased by 1.2%. The soluble sugar content in *L. formosana*, *P. massoniana*, and *P. zhennan* decreased at all the pH levels except pH 5.5, where it showed an increasing trend. The soluble protein content in all the species showed a decreasing trend with the increase in the acidity level (Table 4). The proline content in all the species was greater at pH 2.5, compared with all the other pH levels and the control. In *C. lanceolata*, the proline content increased at pH 2.5 but decreased by 24 and 28% at pH 4.5 and 5.5, respectively. Similarly, in *S. superba*, the proline



Fig. 3. Enzymatic antioxidants and lipid peroxidation under acid deposition treatments. Vertical bars indicate the standard error, the results are different at $p \le 0.05$: a) CAT, b) POD, c) SOD, and d) MDA.

| Tree parameter | Treatment | C. lanceolata | L. formosana | P. masooniana | P. zhennan | S. superba |
|--|-----------|-------------------|--------------|-------------------|--------------------|--------------------|
| Soluble sugar (mg g ⁻¹ FW) | СК | 5.83±1.07a | 10.35±0.66a | 5.34±1.29a | 12.25±0.95a | 4.99±1.11b |
| | 2.5 | 3.25±0.28b | 5.98±0.73b | 2.96±0.20c | 9.67±0.72b | 4.25±0.84b |
| | 3.5 | 3.08±0.31b | 6.59±0.18b | 3.66±0.23b | 10.18±1.03b | 4.94±0.89b |
| | 4.5 | $4.01 \pm 0.02b$ | 4.53±0.31b | 3.12±0.19b | 10.35±0.77b | 5.05±1.05a |
| | 5.5 | $3.22 \pm 0.07 b$ | 6.83±0.73b | 3.67±0.17b | 11.63±0.45a | 4.54±1.09b |
| | | | | | | |
| Protein (mg g ⁻¹ FW) | CK | 4.92±0.06a | 7.94±0.51a | 7.17±0.14a | 9.28±0.43a | 8.69±0.07a |
| | 2.5 | 1.84±0.03c | 1.73±0.02c | 2.55±0.06b | 2.28±0.06c | 2.06±0.05c |
| | 3.5 | 2.07±0.06c | 2.47±0.05c | 2.68±0.21b | 3.20±0.17b | 3.24±0.23b |
| | 4.5 | $2.24{\pm}0.03b$ | 3.84±0.03b | 3.33±0.34b | 3.70±0.13b | 3.25±0.39b |
| | 5.5 | $3.85{\pm}0.08b$ | 4.47±0.15b | 4.92±0.06a | 4.80±0.34a | 3.89±0.34b |
| | | | | | | |
| Proline (mg g ⁻¹ FW) | CK | $0.25 \pm 0.03 b$ | 0.64±0.21b | $0.19{\pm}0.01b$ | 0.15±0.004b | $0.22 \pm 0.007 b$ |
| | 2.5 | 0.59±0.04a | 1.00±0.23a | 0.25±0.03b | $0.16 \pm 0.008 b$ | $0.24 \pm 0.009 b$ |
| | 3.5 | 0.33±0.04b | 0.75±0.17b | $0.19{\pm}0.009b$ | $0.17 \pm 0.008 b$ | 0.36±0.04a |
| | 4.5 | $0.19{\pm}0.01b$ | 0.46±0.01b | $0.24 \pm 0.02b$ | $0.20{\pm}0.008a$ | $0.21 \pm 0.008 b$ |
| | 5.5 | $0.18{\pm}0.005b$ | 0.52±0.02b | 0.24±0.03b | 0.21±0.005a | 0.18±0.02c |

Table 4. Total soluble sugar, protein, and proline content in response to acid stress (mean \pm SE)

content increased by 9.9 and 63.6% at pH 2.5 and pH 3.5 but decreased by 4.5 and 18.1% at pH 4.5 and pH 5.5, as presented in Table 4.

4. DISCUSSION

The present study demonstrated that acid deposition exerts significant detrimental effects on plant growth, as evidenced by reductions in plant height, root collar diameter, and overall biomass. These findings are consistent with previous research by Tong and Liang (2005), who reported pronounced structural damage and visible foliar chlorosis in response to acid deposition at pH levels below 3.5. Such symptoms suggest that acid stress not only impairs aboveground growth but also disrupts physiological processes vital for plant development. Our findings are corroborated with earlier reports mentioning leaf damage in seedlings exposed to pH 3.0 for ten days (Eguagie et al., 2016), marginal necrosis (Bussotti et al., 1997), and curtailed photosynthesis (Flagler et al., 1994). All these symptoms collectively impaired plant height, stem growth, and tree biomass (Song et al., 2011). These cumulative findings strongly reinforce our conclusion that acid deposition at or below pH 3.5 significantly inhibits seedling growth and biomass accumulation. The severity of this inhibition suggests that seedlings exposed to such stress conditions may adopt specific tolerance strategies to mitigate damage. We argue that one primary strategy involves the selective abscission of damaged leaf regions, allowing the plant to preserve healthier tissue and limit further injury. A secondary mechanism may include the formation of protective scars on leaves, which could serve to isolate and contain the damage. Furthermore, our results revealed marked variation in root-to-shoot ratios across different pH treatments, indicating adaptive allocation of resources under stress. This trend aligns with earlier findings in spruce seedlings, where similar changes in biomass partitioning were observed under acid stress, supporting the notion of a broader, conserved physiological response among tree species (Bäck *et al.*, 1995).

4.1. Gas exchange measurements

Photosynthetic rate (P_n) can be influenced by both stomatal and non-stomatal limitations. Stomatal constraints primarily involve the regulation of gas exchange through the opening and closing of stomata, whereas non-stomatal factors affect photosynthesis by altering mesophyll cell activity (Wang et al., 2014). Photosynthesis generates several organic compounds taking critical part in physiological function of plants (Xu et al., 2022). It was observed that the P_n value was reduced with a rise in acidity, which is consistent with our results for C. lanceolata and P. massoniana (Sun et al., 2016). One plausible explanation for this decline is the leaching of magnesium or hydrogen ions from chlorophyll, which may impair chloroplast function and hinder the synthesis of adenosine triphosphate (ATP), a key energy molecule in the photosynthetic process (Verbruggen and Hermans, 2013). The study found that L. formosana, P. zhennan, and S. superba maintained higher photosynthetic rates under acid stress, compared to C. lanceolata and P. massoniana, suggesting greater stomatal tolerance. Additionally, both stomatal and non-stomatal factors, especially enzyme activity, contribute to regulating photosynthesis in acidic conditions (Zhang et al., 2021b). It is well agreed that stomata are vanguards in the process of photosynthesis and particularly influence P_n when G_s and $C_{\rm i}$ are lower. The decrease in $G_{\rm s}$ is in inverse relationship with C_i, which also reflects appropriate and essential dominance of non-stomatal factors (Velikova et al., 1999). Our recorded data revealed that increased C_i reduced G_s in all the species and alleviated acid deposition by reducing transpiration loss. The difference in photosynthetic attributes after acid exposure can also be due to biological and morphological characteristics of species and their resistance to acid deposition (Zhang et al., 2023). Plant species respond differentially to stress, which is a crucial factor in influencing the photosynthetic activity of any plant (Ahmad et al., 2019). The G_s and T_r of S. superba, P. zhennan, and L. formosana were higher at pH 5.5, compared to the control group, revealing that mild stress enhanced the photosynthetic rate in these species. Acid deposition below pH 3.5 had harmful impacts on the photosynthesis activity, and our results agree with those obtained by Shu et al. (2019), who reported the same findings.

4.2. Estimation of biochemical attributes and photosynthetic pigments

Chlorophyll can be used as a sign of resistance to depict the eco physiological condition of plants (Zhou *et al.*, 2020). This study showed that the chlorophyll a, b, and carotenoid content in the five species were lower under different acid levels. Our study agrees with the findings of the study by Moharekar *et al.* (2003), who observed reduction in the chlorophyll content due to a rise in acid deposition stress. The chlorophyll content is frequently affected by strong acid rain (Kumaravelu and Ramanujam, 1998). The main cause of the decline in chlorophyll content might be the foliar leaching of nutrients due to acidity (Bussotti *et al.*, 1997).

The findings reported by Qiu and Liu (2002) demonstrated that elevated acidity levels (pH 2.5 and 3.5) led to a reduction in chlorophyll a content in *Dimocarpus longan* leaves, while chlorophyll *b* was comparatively less affected – an observation that aligns with our results. Similarly, other findings (Jian-Fu *et al.*, 2013) have also confirmed chlorophyll reduction by acid deposition. Carotenoids can protect chlorophylls by scavenging ROS through xanthophyll (Khalid *et al.*, 2018). Our findings revealed that the carotenoid content decreased below pH 3.5 and pH 2.5, compared to that in the control treatment plants; this is perhaps due to degradation of pigments (Zhang *et al.*, 2020). Antioxidant enzymes are produced by plants to deal with stress stimulated by abiotic and biotic stresses. We revealed that, contrary to the control plants, the acid deposition enhanced CAT and POD activities, which is in accordance with a study conducted by Wu *et al.* (2012).

The study found that the increased acid concentration elevated leaf POD activity, indicating heightened plant stress. While P. zhennan and S. superba managed hydrogen peroxide more effectively, L. formosana showed reduced CAT, POD, and SOD activity, supporting earlier findings that it is more sensitive to acid stress (Chen et al., 2013). The initial increase followed by a decrease in enzyme activities observed in this study was attributed to prolonged exposure to acidity. Higher acidity impaired the plants' detoxification capacity, disrupting the balance between protective enzymes and antioxidant scavenging. MDA, an indicator of oxidative damage, significantly increased under the acid treatment, rising with the acidity levels. This trend suggests that acid deposition caused membrane damage, leading to the accumulation of reactive oxygen species (ROS) (Liu et al., 2019). In our trial, MDA in both P. zhennan and S. superba was lower than in the other species, suggesting that these species displayed advanced antioxidant enzyme activities, mitigating the damage caused by ROS, compared to C. lanceolata, P. massoniana, and L. formosana. These findings are in agreement with those shown by Wyrwicka and Skłodowska (2006), Rajjou et al. (2012) and Chen et al. (2013), who observed the same results. These findings suggested that free radical production is enhanced by acid deposition and causes lipid peroxidation of membranes, resulting in increased MDA. This increase exhibits membrane injury, ROS activity, and oxidative stress (Montillet et al., 2005). Plants with improved enzyme activity can defend their cells from ecological pressure (Noman et al., 2015). It is most likely that the antioxidant compounds keep the defensive pathway active, and these seedlings produce enzymes under adapting stress. Soluble sugars are the primary osmoregulators which are significant in sugar detection in plants (Akhter et al., 2022). The soluble sugar of the five species decreased under the acid deposition at all the pH levels. These sugars work as osmoprotectants in maintaining the pressure by stabilizing the cell membrane (Marcińska et al., 2013). In the present study, P. zhennan was least affected by the acid deposition, compared to the other species. The increased sugar level can be viewed as an effective approach of P. zhennan to stop the damaging impact of acid stress (Foyer and Noctor, 2005). It can be concluded that acid deposition increases the demand for nutrients by lowering sugar synthesis (Bao et al., 2019).

Soluble protein is a prominent osmotic plant regulator, as its increased content aids in balancing cells and its resistance to the damage caused by stress (Amini and Ehsanpour, 2005). The results showed a significant reduction in protein content under the acid stress (\leq pH 3.5), compared to the control treatment. A similar study on *Dimocarpus longan* found that stress at \leq pH 3.5 also impacted leaf protein levels, indicating that acid deposition reduces primary respiratory pathways and plant energy (Pan *et al.*, 2015).

Similarly, if the acid stress is high, the activity of enzymes involved in the synthesis of plant proteins is reduced (Bao et al., 2019). Proline acts as an osmolyte that can protect cell dehydration as well a defensive agent and a (ROS) scavenger (Debnath et al., 2018). It increased under the acid stress in the five species. This indicates that proline was involved in osmotic regulation. A previous study also showed that acid stress prevents proline production in Hypogymnia (Kováčik et al., 2011). Our result agrees with those reported by Chen et al. (2013), who found that the proline level in L. formosana, P. zhennan, and S. superba increased under acid stress. A similar study also noted that the stress led to higher proline contents in P. zhennan leaves (Hu et al., 2015). Plants maintain cell shape and protect membranes by generating proline under high acidity, preventing lipid peroxidation and oxidative stress (Zhang et al., 2021a). Our findings are consistent with those obtained by Zhang et al. (2020), who examined tea trees. The observed changes in acid resistance among the species highlight the importance of developing strategies to mitigate damage, which is vital for the sustainability of forest ecosystems.

5. CONCLUSIONS

The present trial concludes that P. massoniana, C. lanceolata, P. zhennan, L. formosana, and S. superba were sensitive to acid deposition, showing varying response among different parameters. Seedling height, root collar diameter, and plant biomass decreased under the acid stress. The levels of chlorophyll a and chlorophyll b were lower than in the control. Seedlings used a protective mechanism to respond differently to stress under osmotic adjustment. The acid deposition significantly increased the antioxidant enzyme activities and aggravated membrane lipid peroxidation. This trial showed improved understanding of photosynthetic, antioxidant, and physiological tolerance of seedlings to numerous kinds of acid deposition. The current findings are useful for the estimation of plant sensitivity to stress and can be valuable for examining plants in acid stress conditions under changing climate. Moreover, the present study aids in identifying P. zhennan, and S. superba as bioindicators and resilient species, compared to the others; hence, extensive planting programs can be beneficial in urban and forest areas to clean the atmosphere and improve the quality of the air.

Conflict of interest. Authors declare no conflict of interest exists.

Author Contributions: Matoor Mohsin Gilani and Bo Liu conceived the idea; Matoor Mohsin Gilani and Xiangqing Ma designed the study. Matoor Mohsin Gilani, Muhammad Haroon U. Rashid and Muhammad Waqqas Khan Tarin performed the lab work, Matoor Mohsin Gilani and Ghulam Yasin analyzed the data. Muhammad Zubair, and Ali Noman discussed the results and discussion. Taimoor Hassan Farooq and Fahad Mohammed Alzuaibr revised the English language.

6. REFERENCES

- Ahmad, I., Gilani, M., Wu, P., 2019. Cuttings growth response of Dalbergia sissoo (shisham) to soil compaction stress. Appl. Ecol. Env. Res. 17(1), 1049-1059. http://dx.doi. org/10.15666/aeer/1701 10491059
- Ahmed, I.M., Cao, F., Zhang, M., Chen, X., Zhang, G., Wu, F., 2013. Difference in yield and physiological features in response to drought and salinity combined stress during anthesis in Tibetan wild and cultivated barleys. PloS one, 8(10) e77869. https://doi.org/10.1371/journal.pone.0077869
- Akhter, N., Habiba, O., Hina, M., Shahnaz, M.M., Alzuaibr, F.M., Alamri, S., et al., 2022. Structural, biochemical, and physiological adjustments for toxicity management, accumulation, and remediation of cadmium in wetland ecosystems by *Typha domingensis* Pers. Water Air Soil Poll. 233(5), 151. https://doi.org/10.1007/s11270-022-05613-w
- Amini, F., Ehsanpour, A.A., 2005. Soluble proteins, proline, carbohydrates and Na+/K+ changes in two tomato (*Lycopersicon esculentum* Mill.) cultivars under *in vitro* salt stress. Am. J. Biochem. Biotech. 1 (4), 204-208. https://doi. org/10.3844/ajbbsp.2005.204.208
- Bäck, J., Huttunen, S., Turunen, M., Lamppu, J., 1995. Effects of acid rain on growth and nutrient concentrations in Scots pine and Norway spruce seedlings grown in a nutrient-rich soil. Environ. Pollut. 89(2), 177-187. https://doi. org/10.1016/0269-7491(94)00054-H
- Bao, G., Tang, W., He, F., Chen, W., Zhu, Y., Fan, C., et al., 2019. Physiological response in the leaf and stolon of white clover under acid precipitation and freeze-thaw stress. Funct. Plant Biol. 47, 50-57. https://doi.org/10.1071/FP19072
- Bates, L.S., Waldren, R., Teare, I., 1973. Rapid determination of free proline for water-stress studies. Plant Soil 39, 205-207. https://doi.org/10.1007/BF00018060
- Blanco, J.A., Wei, X., Jiang, H., Jie, C.Y., Xin, Z.H., 2012. Impacts of enhanced nitrogen deposition and soil acidification on biomass production and nitrogen leaching in Chinese fir plantations. Can. J. For. Res. 42, 437-450. https://doi.org/10.1139/x2012-004
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem. 72, 248-254. https://doi.org/10.1016/0003-2697(76)90527-3
- Bussotti, F., Bottacci, A., Grossoni, P., Mori, B., Tani, C., 1997. Cytological and structural changes in *Pinus pines* L. needles following the application of an anionic surfactant. Plant Cell Environ. 20, 513-520. https://doi. org/10.1046/j.1365-3040.1997.d01-91.x
- Cape, J., 1993. Direct damage to vegetation caused by acid rain and polluted cloud: definition of critical levels for forest trees. Environ. Pollut. 82, 167-180. https://doi. org/10.1016/0269-7491(93)90114-4
- Chang, C.-T., Yang, C.-J., Huang, K.-H., Huang, J.-C., Lin, T.-C., 2022. Changes of precipitation acidity related to sulfur and nitrogen deposition in forests across three continents in north hemisphere over last two decades. Sci. Total Environ. 806, 150552. https://doi.org/10.1016/j.scitotenv.2021.150552

- Chen, J., Hu, W.-J., Wang, C., Liu, T.-W., Chalifour, A., Chen, J., et al., 2014. Proteomic analysis reveals differences in tolerance to acid rain in two broad-leaf tree species, *Liquidambar* formosana and Schima superba. PLoS One, 9(7), e102532. https://doi.org/10.1371/journal.pone.0102532
- Chen, J., Li, W., Gao, F., 2010. Biogeochemical effects of forest vegetation on acid precipitation-related water chemistry: a case study in southwest China. J. Environ. Monit. 12(10), 1799-1806. https://doi.org/10.1039/C0EM00116C
- Chen, J., Wang, W.-H., Liu, T.-W., Wu, F.-H., Zheng, H.-L., 2013. Photosynthetic and antioxidant responses of *Liquidambar formosana* and *Schima superba* seedlings to sulfuric-rich and nitric-rich simulated acid rain. Plant Physiol. Biochem. 64, 41-51. https://doi.org/10.1016/j.plaphy.2012.12.012
- Chen, L., Heerink, N., Van Den Berg, M., 2006. Energy consumption in rural China: A household model for three villages in Jiangxi Province. Ecol. Econ. 58, 407-420. https://doi. org/10.1016/j.ecolecon.2005.07.018
- Chen, Z., Ye, S., Cao, J., Shang, H., 2021. Nitrogen fertilization modified the responses of *Schima superba* seedlings to elevated CO₂ in subtropical China. Plants 10(2), 383. https:// doi.org/10.3390/plants10020383
- Cheng, Y., Peng, J., Gu, Y., Guo, H., Jiang, T., Yang, H., 2023. The early effect of plant density on soil physicochemical attributes and bacterial and understory plant diversity in *Phoebe zhennan* plantations. Forests 14. https://doi.org/10.3390/ f14081612
- Debnath, B., Irshad, M., Mitra, S., Li, M., Rizwan, H.M., Liu, S., et al., 2018. Acid rain deposition modulates photosynthesis, enzymatic and non-enzymatic antioxidant activities in tomato. Int. J. Environ. Res. 12, 203-214. https://doi. org/10.1007/s41742-018-0084-0
- Debnath, B., Sikdar, A., Islam, S., Hasan, K., Li, M., Qiu, D., 2021. Physiological and molecular responses to acid rain stress in plants and the impact of melatonin, glutathione and silicon in the amendment of plant acid rain stress. Molecules 26(4), 862. https://doi.org/10.3390/molecules26040862
- Diatta, J., Youssef, N., Tylman, O., Grzebisz, W., Markert, B., Drobek, L., *et al.*, 2021. Acid rain induced leakage of Ca, Mg, Zn, Fe from plant photosynthetic organs-Testing for deciduous and dicotyledons. Ecol. Indic. 121, 107210. https://doi.org/10.1016/j.ecolind.2020.107210
- Dong, D., Du, E., Sun, Z., Zeng, X., De Vries, W., 2017. Nonlinear direct effects of acid rain on leaf photosynthetic rate of terrestrial plants. Environ. Pollut. 231, 1442-1445. https://doi.org/10.1016/j.envpol.2017.09.005
- Du, E., Dong, D., Zeng, X., Sun, Z., Jiang, X., De Vries, W., 2017. Direct effect of acid rain on leaf chlorophyll content of terrestrial plants in China. Sci. Total Environ. 605, 764-769. https://doi.org/10.1016/j.scitotenv.2017.06.044
- Eguagie, M., Aiwansoba, R., Omofomwan, K., Oyanoghafo, O., 2016. Impact of simulated acid rain on the growth, yield and plant component of *Abelmoschus caillei*. J. Adv. Biol. Biotechnol. 6, 1-6. https://doi.org/0.9734/JABB/2016/24804
- Fan, H.B., Wang, Y.H., 2000. Effects of simulated acid rain on germination, foliar damage, chlorophyll contents and seedling growth of five hardwood species growing in China. For. Ecol. Manag. 126(3), 321-329. https://doi.org/10.1016/ S0378-1127(99)00103-6

- Farooq, T., Yan, W., Rashid, M., Tigabu, M., Gilani, M., Zou, X., Wu, P., 2019. Chinese fir (*Cunninghamia Lanceolata*) a green gold of china with continues decline in its productivity over the successive rotations: A review. Appl. Ecol. Env. Res. 17, 5. https://doi.org/10.15666/acer/1705 1105511067
- Farooq, T.H., Shakoor, A., Wu, X., Li, Y., Rashid, M.H.U., Zhang, X., et al., 2021. Perspectives of plantation forests in the sustainable forest development of China. Iforest. 14, 166-174. http://doi.org/10.3832/ifor3551-014
- Farooq, T.H., Yan, W., Chen, X., Shakoor, A., Rashid, M.H.U., Gilani, M.M., *et al.*, 2020. Dynamics of canopy development of *Cunninghamia lanceolata* mid-age plantation in relation to foliar nitrogen and soil quality influenced by stand density. Glob. Ecol. Conserv. 24, e01209. https://doi. org/10.1016/j.gecco.2020.e01209
- Feng, Z.-W., Miao, H., Zhang, F.-Z., Huang, Y.-Z., 2002. Effects of acid deposition on terrestrial ecosystems and their rehabilitation strategies in China. J. Environ Sci. (China) 14(2), 227-33. PMID: 12046292.
- Flagler, R.B., Lock, J.E., Elsik, C.G., 1994. Leaf-level and wholeplant gas exchange characteristics of shortleaf pine exposed to ozone and simulated acid rain. Tree Physiol. 14, 361-374. https://doi.org/10.1093/treephys/14.4.361
- Foyer, C.H., Noctor, G., 2005. Oxidant and antioxidant signalling in plants: a re-evaluation of the concept of oxidative stress in a physiological context. Plant Cell Environ. 28, 1056-1071. https://doi.org/10.1111/j.1365-3040.2005.01327.x
- Gilani, M., Tigabu, M., Farooq, T., Rashid, M., Yasin, G., Zubair, M., et al., 2023. Influence of differential nitrogen forms on seedling morphology of four widely grown species in southern China. Appl. Ecol. Env. Res. 21. https://doi. org/10.15666/aeer/2106 59896002
- Gilani, M.M., Tigabu, M., Liu, B., Farooq, T.H., Rashid, M.H.U., Ramzan, M., *et al.*, 2021. Seed germination and seedling emergence of four tree species of southern China in response to acid rain. J. For. Res. 32, 471-481. https://doi. org/10.1007/s11676-020-01102-0
- Grennfelt, P., Engleryd, A., Forsius, M., Hov, Ø., Rodhe, H., Cowling, E., 2020. Acid rain and air pollution: 50 years of progress in environmental science and policy. Ambio 49, 849-864. https://doi.org/10.1007/s13280-019-01244-4
- Hasan, M.M., Liu, X.-D., Waseem, M., Guang-Qian, Y., Alabdallah, N.M., Jahan, M.S., *et al.*, 2022. ABA activated SnRK2 kinases: An emerging role in plant growth and physiology. Plant Signal Behav. 17(1), 2071024. https:// doi.org/10.1080/15592324.2022.2071024
- Hodges, D.M., Delong, J.M., Forney, C.F., Prange, R.K., 1999. Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. Planta 207, 604-611. https://doi.org/10.1007/s004250050524
- Hu, W.-J., Chen, J., Liu, T.-W., Simon, M., Wang, W.-H., Wu, F.-H., et al., 2014. Comparative proteomic analysis of differential responses of Pinus massoniana and *Taxus* wallichiana var. mairei to simulated acid rain. Int. J. Mol. Sci. 15(3), 4333-4355. https://doi.org/10.3390/ijms15034333
- Hu, Y., Wang, B., Hu, T., Chen, H., Li, H., Zhang, W., et al., 2015. Combined action of an antioxidant defence system and

osmolytes on drought tolerance and post-drought recovery of *Phoebe zhennan S. Lee* saplings. Acta Physiol. Plant. 37, 1-13. https://doi.org/10.1007/s11738-015-1831-x

- Hua, T., Tan, S., Liu, Z., 2022. Effects of cadmium and simulated acid rain on growth and physiological characteristics of bermudagrass seedling. Acta Physiol. Plant. 44, 17. https:// doi.org/10.1007/s11738-021-03353-8
- Huang, L., Yang, J., Zhang, G., 2012. Chemistry and source identification of wet precipitation in a rural watershed of subtropical China. Chin. J. Geochem. 31, 347-354. https:// doi.org/10.1007/s11631-012-0585-z
- Jian-Fu, L., Ming-Yuan, W., Chen, Y., Ai-Jun, Z., 2013. Effects of exogenous nitric oxide on physiological characteristics of longan (*Dimocarpus longana*) seedlings under acid rain stress. Yingyong Shengtai Xuebao 24(8). https://openurl. ebsco.com/EPDB%3Agcd%3A5%3A25872389/detailv2?s id=ebsco%3Aplink%3Ascholar&id=ebsco%3Agcd%3A98 381864&crl=c&link origin=scholar.google.com
- Khalid, N., Noman, A., Aqeel, M., Hadayat, N., Anjum, S., 2018. NPK could alleviate the adverse effects of simulated acid rain in sunflower (*Helianthus annuus* L.). Plant Nutr. 41(5), 584-595. https://doi.org/10.1080/01904167.2017.1406108
- Kováčik, J., Klejdus, B., Bačkor, M., Štork, F., Hedbavny, J., 2011. Physiological responses of root-less epiphytic plants to acid rain. Ecotoxicol. 20, 348-357. https://doi. org/10.1007/s10646-010-0585-x
- Kumaravelu, G., Ramanujam, M., 1998. Impact of simulated acidic rain on growth, photosynthetic pigments, cell metabolites, and leaf characteristics of green gram. Photosynthetica 35, 71-78. https://doi.org/10.1023/A:1006821931577
- Li, Z., Peng, Y., Ma, X., 2013. Different response on drought tolerance and post-drought recovery between the small-leafed and the large-leafed white clover (*Trifolium repens* L.) associated with antioxidative enzyme protection and lignin metabolism. Acta Physiol. Plant. 35, 213-222. https://doi. org/10.1007/s11738-012-1066-z
- Lichtenthaler, H.K., Wellburn, A.R., 1983. Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. Portland Press Ltd. https://doi. org/10.1042/bst0110591
- Liu, J., Zhou, G., Yang, C., Ou, Z., Peng, C., 2007. Responses of chlorophyll fluorescence and xanthophyll cycle in leaves of *Schima superba* Gardn. Champ. and *Pinus massoniana* Lamb. to simulated acid rain at Dinghushan Biosphere Reserve, China. Acta Physiol. Plant. 29, 33-38. https://doi. org/10.1007/s11738-006-0005-2
- Liu, M., Korpelainen, H., Dong, L., Yi, L., 2019. Physiological responses of *Elaeocarpus glabripetalus* seedlings exposed to simulated acid rain and cadmium. Ecotoxicol. Environ. Saf. 175, 118-127. https://doi.org/10.1016/j.ecoenv.2019.03.026
- Liu, X., Fu, Z., Zhang, B., Zhai, L., Meng, M., Lin, J., et al., 2018. Effects of sulfuric, nitric, and mixed acid rain on Chinese fir sapling growth in Southern China. Ecotoxicol. Environ. Saf. 160, 154-161. https://doi.org/10.1016/j.ecoenv.2018.04.071
- Liu, Z., Liu, Z., Wu, L., Li, Y., Wang, J., Wei, H., et al., 2022. Effect of polyethylene microplastics and acid rain on the agricultural soil ecosystem in Southern China. Environ. Pollut. 303, 119094. https://doi.org/10.1016/j.envpol.2022.119094
- Ma, S., Chen, W., Zhang, J., Shen, H., 2020. Influence of simulated acid rain on the physiological response of flowering Chinese cabbage and variation of soil nutrients. Plant Soil Environ. 66. https://doi.org/10.17221/469/2020-PSE

- Marcińska, I., Czyczyło-Mysza, I., Skrzypek, E., Grzesiak, M.T., Janowiak, F., Filek, M., *et al.*, 2013. Alleviation of osmotic stress effects by exogenous application of salicylic or abscisic acid on wheat seedlings. Int. J. Mol. Sci. 14, 13171-13193. https://doi.org/10.3390/ijms140713171
- Meng, Y., Zhao, Y., Li, R., Li, J., Cui, L., Kong, L., et al., 2019. Characterization of inorganic ions in rainwater in the megacity of Shanghai: Spatiotemporal variations and source apportionment. Atmos. Res. 222, 12-24. https://doi. org/10.1016/j.atmosres.2019.01.023
- Mofunanya, A., Soonen, L., 2017. Physiological and morphological responses of *Amaranthus hybridus* L. (green) to simulated nitric and sulphuric acid rain. Br. J. Appl. Sci. Technol. 21, 1-12. https://doi.org/10.9734/BJAST/2017/31863
- Moharekar, S., Lokhande, S., Hara, T., Tanaka, R., Tanaka, A., Chavan, P., 2003. Effect of salicylic acid on chlorophyll and carotenoid contents of wheat and moong seedlings. Photosynthetica 41, 315-317. https://doi. org/10.1023/B:PHOT.0000011970.62172.15
- Montillet, J.-L., Chamnongpol, S., Rustérucci, C., Dat, J., Van De Cotte, B., Agnel, J.-P., *et al.*, 2005. Fatty acid hydroperoxides and H₂O₂ in the execution of hypersensitive cell death in tobacco leaves. Plant Physiol. 138, 1516-1526. https:// doi.org/10.1104/pp.105.059907
- Naheed, R., Aslam, H., Kanwal, H., Farhat, F., Gamar, M.I.A., Al-Mushhin, A.A., *et al.*, 2021. Growth attributes, biochemical modulations, antioxidant enzymatic metabolism and yield in *Brassica napus* varieties for salinity tolerance. Saudi J. Biol. Sci. 28(10), 5469-5479. https://doi. org/10.1016/j.sjbs.2021.08.021
- Noman, A., Ali, Q., Maqsood, J., Iqbal, N., Javed, M.T., Rasool, N., et al., 2018. Deciphering physio-biochemical, yield, and nutritional quality attributes of water-stressed radish (*Raphanus sativus* L.) plants grown from Zn-Lys primed seeds. Chemosphere 195, pp.175-189. https://doi. org/10.1016/j.chemosphere.2017.12.059
- Noman, A., Ali, S., Naheed, F., Ali, Q., Farid, M., Rizwan, M., et al., 2015. Foliar application of ascorbate enhances the physiological and biochemical attributes of maize (Zea mays L.) cultivars under drought stress. Archives Agron. Soil Sci. 61(12), 1659-1672. https://doi.org/10.1080/03650 340.2015.1028379
- Pan, T., Li, Y., Ma, C., Qiu, D., 2015. Calcium affecting protein expression in longan under simulated acid rain stress. ESPR 22, 12215-12223. https://doi.org/10.1007/s11356-015-4389-7
- Qiu, D., Liu, X., 2002. Effects of simulated acid rain on chloroplast activity in *Dimorcarpus longana* Lour. *cv.* wulongling leaves. Ying Yong Sheg Tai Xue Bao. J. Appl. Ecol. 13, 1559-1562. https://europepmc.org/article/med/12682955
- Rahman, S.U., Xuebin, Q., Kamran, M., Yasin, G., Cheng, H., Rehim, A., *et al.*, 2021. Silicon elevated cadmium tolerance in wheat (*Triticum aestivum* L.) by endorsing nutrients uptake and antioxidative defense mechanisms in the leaves. Plant Physiol. Biochem. 166, 148-159. https://doi. org/10.1016/j.plaphy.2021.05.038
- Rajjou, L., Duval, M., Gallardo, K., Catusse, J., Bally, J., Job, C., et al., 2012. Seed germination and vigor. Annu. Rev. Plant Biol. 63, 507-533. https://doi.org/10.1146/ annurev-arplant-042811-105550
- Ren, J., Zhu, L., Zhang, X., Luo, Y., Zhong, X., Li, B., *et al.*, 2024. Variation characteristics of acid rain in Zhuzhou, Central China over the period 2011-2020. J. Environ. Sci. 138, 496-505. https://doi.org/10.1016/j.jes.2023.03.035

- Shu, X., Zhang, K., Zhang, Q., Wang, W., 2019. Ecophysiological responses of *Jatropha curcas* L. seedlings to simulated acid rain under different soil types. Ecotoxicol. Environ. Saf. 185, 109705. https://doi.org/10.1016/j.ecoenv.2019.109705
- Song, Y., Li, D., Wang, Y., Wang, R., Xu, H., 2011. Chemical characteristics and cause analysis of precipitation in the South of Liaodong Peninsula. Procedia Eng. 18, 220-225. https://doi.org/10.1016/j.proeng.2011.11.035
- Sun, J., Hu, H., Li, Y., Wang, L., Zhou, Q., Huang, X., 2016. Effects and mechanism of acid rain on plant chloroplast ATP synthase. ESPR. 23, 18296-18306. https://doi. org/10.1007/s11356-016-7016-3
- Sun, Z., Wang, L., Chen, M., Wang, L., Liang, C., Zhou, Q., et al., 2012. Interactive effects of cadmium and acid rain on photosynthetic light reaction in soybean seedlings. Ecotoxicol. Environ. Saf. 79, 62-68. https://doi.org/10.1016/j. ecoenv.2011.12.004
- Tong, G., Liang, H., 2005. Effects of simulated acid rain and its acidified soil on soluble sugar and nitrogen contents of wheat seedlings (in China). Ying Yong Sheng Tai Xue Bao 16(8), 1487-92. Chinese. PMID: 16262064. https://pubmed.ncbi.nlm.nih.gov/16262064/
- Velikova, V., Tsonev, T., Yordanov, I., 1999. Light and CO₂ responses of photosynthesis and chlorophyll fluorescence characteristics in bean plants after simulated acid rain. Physiol Plant. 107, 77-83. https://doi. org/10.1034/j.1399-3054.1999.100111.x
- Verbruggen, N., Hermans, C., 2013. Physiological and molecular responses to magnesium nutritional imbalance in plants. Plant Soil 368, 87-99. https://doi.org/10.1007/s11104-013-1589-0
- Wang, L., Wang, W., Zhou, Q., Huang, X., 2014. Combined effects of lanthanum (III) chloride and acid rain on photosynthetic parameters in rice. Chemosphere 112, 355-361. https://doi.org/10.1016/j.chemosphere.2014.04.069
- Wei, H., Liu, W., Zhang, J., Qin, Z., 2017. Effects of simulated acid rain on soil fauna community composition and their ecological niches. Environ. Pollut. 220, 460-468. https:// doi.org/10.1016/j.envpol.2016.09.088
- Wu Gang, W. G., 1998. Effect of acidic deposition on productivity of forest ecosystem and estimation of its economic losses in southern suburbs of Chongqing, China 210-215. https:// www.cabidigitallibrary.org/doi/full/10.5555/19980611100
- Wu, H., Wu, X., Li, Z., Duan, L., Zhang, M., 2012. Physiological evaluation of drought stress tolerance and recovery in cauliflower (*Brassica oleracea* L.) seedlings treated with methyl jasmonate and coronatine. J. Plant Growth Regul. 31, 113-123. https://doi.org/10.1007/s00344-011-9224-x
- Wu, J., Wang, M., Wang, T., Fu, X., 2022. Evaluation of ecological service function of liquidambar formosana plantations. Int. J. Environ. Res. Public Health. 19, 15317. https://doi. org/10.3390/ijerph192215317
- Wyrwicka, A., Skłodowska, M., 2006. Influence of repeated acid rain treatment on antioxidative enzyme activities and on lipid peroxidation in cucumber leaves. Environ. Exp. Bot. 56, 198-204. https://doi.org/10.1016/j.envexpbot.2005.02.003

- Xu, J., Guo, L., Liu, L., 2022. Exogenous silicon alleviates drought stress in maize by improving growth, photosynthetic and antioxidant metabolism. Environ. Exp. Bot. 201, 104974. https://doi.org/10.1016/j.envexpbot.2022.104974
- Yao, F.-F., Ding, H.-M., Feng, L.-L., Chen, J.-J., Yang, S.-Y., Wang, X.-H., 2016. Photosynthetic and growth responses of *Schima superba* seedlings to sulfuric and nitric acid depositions. ESPR. 23, 8644-8658. https://doi.org/10.1007/ s11356-015-5970-9
- Yasin, G., Nawaz, M.F., Zubair, M., Azhar, M.F., Gilani, M.M., Ashraf, M.N., *et al.*, 2023. Role of traditional agroforestry systems in climate change mitigation through carbon sequestration: An investigation from the semi-arid region of Pakistan. Land, 12(2), 513. https://doi.org/10.3390/land12020513
- Yasin, G., Ur Rahman, S., Yousaf, M.T.B., Azhar, M.F., Zahid, D.M., Imtiaz, M., *et al.*, 2021. Phytoremediation potential of *E. camaldulensis* and *M. alba* for copper, cadmium, and lead absorption in urban areas of Faisalabad City, Pakistan. International J. Environ. Res. 15, 597-612. https://doi. org/10.1007/s41742-021-00330-4
- Yemm, E., Willis, A., 1954. The estimation of carbohydrates in plant extracts by anthrone. Biochem. J. 57, 508. https://doi. org/10.1042/bj0570508
- Zhang, C., Yi, X., Gao, X., Wang, M., Shao, C., Lv, Z., et al., 2020. Physiological and biochemical responses of tea seedlings (*Camellia sinensis*) to simulated acid rain conditions. Ecotoxicol. Environ. Saf. 192, 110315. https://doi. org/10.1016/j.ecoenv.2020.110315
- Zhang, Y., Tian, C., Yu, T., Dayananda, B., Fu, B., Senaratne, S.L., et al., 2021a. Differential effects of acid rain on photosynthetic performance and pigment composition of the critically endangered Acer amplum subsp. catalpifolium. Glob. Ecol. Conserv. 30, e01773. https://doi.org/10.1016/j. gecco.2021.e01773
- Zhang, Y., Yu, T., Ma, W., Dayananda, B., Iwasaki, K., Li, J., 2021b. Morphological, physiological and photophysiological responses of critically endangered acer catalpifolium to acid stress. Plants 10, 1958. https://doi.org/10.3390/ plants10091958
- Zhang, Y., Yang, F., Wang, Y., Zheng, Y., Zhu, J., 2023. Effects of acid rain stress on the physiological and biochemical characteristics of three plant species. Forests 14, 1067. https:// doi.org/10.3390/f14051067
- Zheng, Q.-P., Wang, H., Chen, B.-B., Sui, P., Lin, W., 2014. Characteristics and the impact factors of acid rain in Fuzhou and Xiamen 1992-2012 (in China). Huan Jing ke Xue Huanjing Kexue, 35, 3644-3650. PMID: 25693365. https:// europepmc.org/article/med/25693365
- Zhou, S., Zhang, M., Chen, S., Xu, W., Zhu, L., Gong, S., et al., 2020. Acid resistance of Masson pine (*Pinus massoniana* Lamb.) families and their root morphology and physiological response to simulated acid deposition. Sci. Rep. 10, 22066. https://doi.org/10.1038/s41598-020-79043-1
- Zhu, Y., An, W., Peng, J., Li, J., Gu, Y., Jiang, B., et al., 2022. Genetic diversity of Nanmu (*Phoebe zhennan S. Lee.* et FN Wei) breeding population and extraction of core collection using nSSR, cpSSR and phenotypic markers. Forests 13, 1320. https://doi.org/10.3390/f13081320